

LAMPIRAN

CORRELATIONS

/VARIABLES=x1.1 x1.2 x1.3 x1.4 x1.5 x1.6 x1.7 x1.8 Kompensasi_X1

/PRINT=TWOTAIL NOSIG

/MISSING=PAIRWISE.

Correlations

Notes

Output Created	01-AUG-2017 12:19:58	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	27
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=x1.1 x1.2 x1.3 x1.4 x1.5 x1.6 x1.7 x1.8 Kompensasi_X1 /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,05
	Elapsed Time	00:00:00,03

Correlations

		x1.1	x1.2	x1.3	x1.4	x1.5
x1.1	Pearson Correlation	1	,612**	,375	,399*	1,000**
	Sig. (2-tailed)		,001	,054	,039	,000
	N	27	27	27	27	27
x1.2	Pearson Correlation	,612**	1	,544**	,627**	,612**
	Sig. (2-tailed)	,001		,003	,000	,001
	N	27	27	27	27	27
x1.3	Pearson Correlation	,375	,544**	1	,490**	,375
	Sig. (2-tailed)	,054	,003		,009	,054
	N	27	27	27	27	27
x1.4	Pearson Correlation	,399*	,627**	,490**	1	,399*
	Sig. (2-tailed)	,039	,000	,009		,039
	N	27	27	27	27	27
x1.5	Pearson Correlation	1,000**	,612**	,375	,399*	1
	Sig. (2-tailed)	,000	,001	,054	,039	
	N	27	27	27	27	27
x1.6	Pearson Correlation	,612**	1,000**	,544**	,627**	,612**
	Sig. (2-tailed)	,001	,000	,003	,000	,001
	N	27	27	27	27	27
x1.7	Pearson Correlation	,375	,544**	1,000**	,490**	,375
	Sig. (2-tailed)	,054	,003	,000	,009	,054
	N	27	27	27	27	27
x1.8	Pearson Correlation	,399*	,627**	,490**	1,000**	,399*
	Sig. (2-tailed)	,039	,000	,009	,000	,039
	N	27	27	27	27	27
Kompensasi_X1	Pearson Correlation	,742**	,858**	,772**	,803**	,742**
	Sig. (2-tailed)	,000	,000	,000	,000	,000
	N	27	27	27	27	27

Correlations

		x1.6	x1.7	x1.8	Kompensasi_X1
x1.1	Pearson Correlation	,612	,375**	,399	,742*
	Sig. (2-tailed)	,001	,054	,039	,000
	N	27	27	27	27
x1.2	Pearson Correlation	1,000**	,544	,627**	,858**
	Sig. (2-tailed)	,000	,003	,000	,000
	N	27	27	27	27
x1.3	Pearson Correlation	,544	1,000**	,490	,772**
	Sig. (2-tailed)	,003	,000	,009	,000
	N	27	27	27	27
x1.4	Pearson Correlation	,627*	,490**	1,000**	,803**
	Sig. (2-tailed)	,000	,009	,000	,000
	N	27	27	27	27
x1.5	Pearson Correlation	,612**	,375**	,399	,742*
	Sig. (2-tailed)	,001	,054	,039	,000
	N	27	27	27	27
x1.6	Pearson Correlation	1**	,544**	,627**	,858**
	Sig. (2-tailed)		,003	,000	,000
	N	27	27	27	27
x1.7	Pearson Correlation	,544	1**	,490**	,772**
	Sig. (2-tailed)	,003		,009	,000
	N	27	27	27	27
x1.8	Pearson Correlation	,627*	,490**	1**	,803**
	Sig. (2-tailed)	,000	,009		,000
	N	27	27	27	27
Kompensasi_X1	Pearson Correlation	,858**	,772**	,803**	1**
	Sig. (2-tailed)	,000	,000	,000	
	N	27	27	27	27

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).

RELIABILITY

/VARIABLES=x1.1 x1.2 x1.3 x1.4 x1.5 x1.6 x1.7 x1.8

/SCALE('ALL VARIABLES') ALL

/MODEL=ALPHA.

Reliability

Notes

Output Created		01-AUG-2017 12:20:16
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	27
Missing Value Handling	Matrix Input	
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.

Syntax		RELIABILITY /VARIABLES=x1.1 x1.2 x1.3 x1.4 x1.5 x1.6 x1.7 x1.8 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01

Scale: ALL VARIABLES

Case Processing Summary

		N	%
Cases	Valid	27	100,0
	Excluded ^a	0	,0
	Total	27	100,0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
,912	8

CORRELATIONS

/VARIABLES=x.2.1 x2.2 x2.3 x2.4 x.2.5 x.2.6 x.2.7 x2.8 x2.9 x2.10 Disiplin.Kerja_X2
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

Correlations

Notes

Output Created		01-AUG-2017 12:22:21
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	27
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=x.2.1 x2.2 x2.3 x2.4 x.2.5 x.2.6 x.2.7 x2.8 x2.9 x2.10 Disiplin.Kerja_X2 /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02

Correlations

		x.2.1	x.2.2	x.2.3	x.2.4	x.2.5
x.2.1	Pearson Correlation	1	,591**	,573**	,684**	1,000**
	Sig. (2-tailed)		,001	,002	,000	,000
	N	27	27	27	27	27
x.2.2	Pearson Correlation	,591**	1	,645**	,647**	,591**
	Sig. (2-tailed)	,001		,000	,000	,001
	N	27	27	27	27	27
x.2.3	Pearson Correlation	,573**	,645**	1	,685**	,573**
	Sig. (2-tailed)	,002	,000		,000	,002
	N	27	27	27	27	27
x.2.4	Pearson Correlation	,684**	,647**	,685**	1	,684**
	Sig. (2-tailed)	,000	,000	,000		,000
	N	27	27	27	27	27
x.2.5	Pearson Correlation	1,000**	,591**	,573**	,684**	1
	Sig. (2-tailed)	,000	,001	,002	,000	
	N	27	27	27	27	27
x.2.6	Pearson Correlation	,654**	,860**	,605**	,640**	,654**
	Sig. (2-tailed)	,000	,000	,001	,000	,000
	N	27	27	27	27	27
x.2.7	Pearson Correlation	,515**	,594**	,936**	,685**	,515**
	Sig. (2-tailed)	,006	,001	,000	,000	,006
	N	27	27	27	27	27
x.2.8	Pearson Correlation	,671**	,721**	,719**	,981**	,671**
	Sig. (2-tailed)	,000	,000	,000	,000	,000
	N	27	27	27	27	27
x.2.9	Pearson Correlation	,597**	,894**	,649**	,606**	,597**
	Sig. (2-tailed)	,001	,000	,000	,001	,001
	N	27	27	27	27	27
x.2.10	Pearson Correlation	,655**	,717**	,969**	,769**	,655**
	Sig. (2-tailed)	,000	,000	,000	,000	,000
	N	27	27	27	27	27
Disiplin.Kerja_X2	Pearson Correlation	,812**	,858**	,848**	,868**	,812**
	Sig. (2-tailed)	,000	,000	,000	,000	,000
	N	27	27	27	27	27

Correlations

		x.2.6	x.2.7	x.2.8	x.2.9	x.2.10
x.2.1	Pearson Correlation	,654	,515**	,671**	,597**	,655**
	Sig. (2-tailed)	,000	,006	,000	,001	,000
	N	27	27	27	27	27
x.2.2	Pearson Correlation	,860**	,594	,721**	,894**	,717**
	Sig. (2-tailed)	,000	,001	,000	,000	,000
	N	27	27	27	27	27
x.2.3	Pearson Correlation	,605**	,936**	,719	,649**	,969**
	Sig. (2-tailed)	,001	,000	,000	,000	,000
	N	27	27	27	27	27
x.2.4	Pearson Correlation	,640**	,685**	,981**	,606	,769**
	Sig. (2-tailed)	,000	,000	,000	,001	,000
	N	27	27	27	27	27
x.2.5	Pearson Correlation	,654**	,515**	,671**	,597**	,655
	Sig. (2-tailed)	,000	,006	,000	,001	,000
	N	27	27	27	27	27
x.2.6	Pearson Correlation	1**	,605**	,668**	,960**	,682**
	Sig. (2-tailed)		,001	,000	,000	,000
	N	27	27	27	27	27
x.2.7	Pearson Correlation	,605**	1**	,669**	,649**	,904**
	Sig. (2-tailed)	,001		,000	,000	,000

	N	27	27	27	27	27
x2.8	Pearson Correlation	,668**	,669**	1**	,638**	,806**
	Sig. (2-tailed)	,000	,000		,000	,000
x2.9	N	27	27	27	27	27
	Pearson Correlation	,960**	,649**	,638**	1**	,725**
x2.10	Sig. (2-tailed)	,000	,000	,000		,000
	N	27	27	27	27	27
Disiplin.Kerja_X2	Pearson Correlation	,682**	,904**	,806**	,725**	1**
	Sig. (2-tailed)	,000	,000	,000	,000	
Disiplin.Kerja_X2	N	27	27	27	27	27
	Pearson Correlation	,866**	,815**	,888**	,862**	,913**
Disiplin.Kerja_X2	Sig. (2-tailed)	,000	,000	,000	,000	,000
	N	27	27	27	27	27

Correlations

		Disiplin.Kerja_X2
x.2.1	Pearson Correlation	,812
	Sig. (2-tailed)	,000
x.2.2	N	27
	Pearson Correlation	,858**
x.2.3	Sig. (2-tailed)	,000
	N	27
x.2.4	Pearson Correlation	,848**
	Sig. (2-tailed)	,000
x.2.5	N	27
	Pearson Correlation	,868**
x.2.6	Sig. (2-tailed)	,000
	N	27
x.2.7	Pearson Correlation	,812**
	Sig. (2-tailed)	,000
x.2.8	N	27
	Pearson Correlation	,866**
x.2.9	Sig. (2-tailed)	,000
	N	27
x.2.10	Pearson Correlation	,815**
	Sig. (2-tailed)	,000
Disiplin.Kerja_X2	N	27
	Pearson Correlation	,888**
Disiplin.Kerja_X2	Sig. (2-tailed)	,000
	N	27
Disiplin.Kerja_X2	Pearson Correlation	,862**
	Sig. (2-tailed)	,000
Disiplin.Kerja_X2	N	27
	Pearson Correlation	,913**
Disiplin.Kerja_X2	Sig. (2-tailed)	,000
	N	27
Disiplin.Kerja_X2	Pearson Correlation	1**
	Sig. (2-tailed)	
Disiplin.Kerja_X2	N	27

** . Correlation is significant at the 0.01 level (2-tailed).

RELIABILITY

/VARIABLES=x.2.1 x2.2 x2.3 x2.4 x.2.5 x.2.6 x.2.7 x2.8 x2.9 x2.10
 /SCALE('ALL VARIABLES') ALL
 /MODEL=ALPHA.

Reliability

Notes

Output Created		01-AUG-2017 12:22:38
Comments		
Input	Active Dataset Filter Weight Split File N of Rows in Working Data File Matrix Input	DataSet0 <none> <none> <none> 27
Missing Value Handling	Definition of Missing Cases Used	User-defined missing values are treated as missing. Statistics are based on all cases with valid data for all variables in the procedure. RELIABILITY /VARIABLES=x.2.1 x2.2 x2.3 x2.4 x.2.5 x.2.6 x.2.7 x2.8 x2.9 x2.10 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA.
Syntax		
Resources	Processor Time Elapsed Time	00:00:00,02 00:00:00,01

Scale: ALL VARIABLES

Case Processing Summary

		N	%
Cases	Valid	27	100,0
	Excluded ^a	0	,0
	Total	27	100,0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
,958	10

CORRELATIONS

/VARIABLES=y.1 y.2 y.3 y.4 y.5 y.6 y.7 y.8 y.9 y.10 Kinerja_Y
 /PRINT=TWOTAIL NOSIG
 /MISSING=PAIRWISE.

Correlations

Notes

Output Created	01-AUG-2017 12:24:43	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	27
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=y.1 y.2 y.3 y.4 y.5 y.6 y.7 y.8 y.9 y.10 Kinerja_Y /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02

Correlations

		y.1	y.2	y.3	y.4	y.5	y.6
y.1	Pearson Correlation	1	,565**	,484*	,493**	,477*	,359
	Sig. (2-tailed)		,002	,010	,009	,012	,066
	N	27	27	27	27	27	27
y.2	Pearson Correlation	,565**	1	,332	,504**	,568**	,489**
	Sig. (2-tailed)	,002		,090	,007	,002	,010
	N	27	27	27	27	27	27
y.3	Pearson Correlation	,484*	,332	1	,718**	,500**	,528**
	Sig. (2-tailed)	,010	,090		,000	,008	,005
	N	27	27	27	27	27	27
y.4	Pearson Correlation	,493**	,504**	,718**	1	,472*	,489**
	Sig. (2-tailed)	,009	,007	,000		,013	,010
	N	27	27	27	27	27	27
y.5	Pearson Correlation	,477*	,568**	,500**	,472*	1	,348
	Sig. (2-tailed)	,012	,002	,008	,013		,075
	N	27	27	27	27	27	27
y.6	Pearson Correlation	,359	,489**	,528**	,489**	,348	1
	Sig. (2-tailed)	,066	,010	,005	,010	,075	
	N	27	27	27	27	27	27
y.7	Pearson Correlation	,493**	,901**	,236	,404*	,472*	,385*
	Sig. (2-tailed)	,009	,000	,236	,036	,013	,047
	N	27	27	27	27	27	27
y.8	Pearson Correlation	,492**	,280	,883**	,711**	,427*	,447*
	Sig. (2-tailed)	,009	,157	,000	,000	,026	,019
	N	27	27	27	27	27	27
y.9	Pearson Correlation	,257	,349	,753**	,914**	,407*	,512**
	Sig. (2-tailed)	,196	,074	,000	,000	,035	,006
	N	27	27	27	27	27	27
y.10	Pearson Correlation	,406*	,568**	,500**	,472*	,906**	,449*
	Sig. (2-tailed)	,035	,002	,008	,013	,000	,019
	N	27	27	27	27	27	27
Kinerja_Y	Pearson Correlation	,701**	,744**	,805**	,833**	,750**	,665**
	Sig. (2-tailed)	,000	,000	,000	,000	,000	,000
	N	27	27	27	27	27	27

Correlations

		y.7	y.8	y.9	y.10	Kinerja_Y
y.1	Pearson Correlation	,493	,492**	,257*	,406**	,701*
	Sig. (2-tailed)	,009	,009	,196	,035	,000
	N	27	27	27	27	27
y.2	Pearson Correlation	,901**	,280	,349	,568**	,744**
	Sig. (2-tailed)	,000	,157	,074	,002	,000
	N	27	27	27	27	27
y.3	Pearson Correlation	,236*	,883	,753	,500**	,805**
	Sig. (2-tailed)	,236	,000	,000	,008	,000
	N	27	27	27	27	27
y.4	Pearson Correlation	,404**	,711**	,914**	,472	,833*
	Sig. (2-tailed)	,036	,000	,000	,013	,000
	N	27	27	27	27	27
y.5	Pearson Correlation	,472*	,427**	,407**	,906*	,750
	Sig. (2-tailed)	,013	,026	,035	,000	,000
	N	27	27	27	27	27
y.6	Pearson Correlation	,385	,447**	,512**	,449**	,665
	Sig. (2-tailed)	,047	,019	,006	,019	,000
	N	27	27	27	27	27
y.7	Pearson Correlation	1**	,194**	,255	,472*	,643*
	Sig. (2-tailed)		,331	,200	,013	,000
	N	27	27	27	27	27
y.8	Pearson Correlation	,194**	1	,732**	,427**	,764*
	Sig. (2-tailed)	,331		,000	,026	,000
	N	27	27	27	27	27
y.9	Pearson Correlation	,255	,732	1**	,407**	,747*
	Sig. (2-tailed)	,200	,000		,035	,000
	N	27	27	27	27	27
y.10	Pearson Correlation	,472*	,427**	,407**	1*	,750**
	Sig. (2-tailed)	,013	,026	,035		,000
	N	27	27	27	27	27
Kinerja_Y	Pearson Correlation	,643**	,764**	,747**	,750**	1**
	Sig. (2-tailed)	,000	,000	,000	,000	
	N	27	27	27	27	27

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).

RELIABILITY

/VARIABLES=y.1 y.2 y.3 y.4 y.5 y.6 y.7 y.8 y.9 y.10

/SCALE('ALL VARIABLES') ALL

/MODEL=ALPHA.

Reliability

Notes

Output Created		01-AUG-2017 12:24:50
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	27
	File	
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.

Syntax	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure. RELIABILITY /VARIABLES=y.1 y.2 y.3 y.4 y.5 y.6 y.7 y.8 y.9 y.10 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,01

Scale: ALL VARIABLES

Case Processing Summary

		N	%
Cases	Valid	27	100,0
	Excluded ^a	0	,0
	Total	27	100,0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
,906	10